DIAGNOSTIC AND THERAPEUTIC CHALLENGES OF VESICOURETERAL REFLUX IN CHILDREN

Sobirjonov Javoxirbek Solijon o‘g‘li
Fergana Premium Medical Clinic Pediatric Urologist

Annotation: *This study examines the diagnostic and therapeutic challenges of vesicoureteral reflux in the pediatric population, analyzing contemporary diagnostic modalities and individualized treatment strategies. The findings emphasize the clinical significance of early detection and risk-stratified intervention, offering important implications for reducing long-term renal morbidity in affected children.*

Keywords: *vesicoureteral reflux, pediatric urology, voiding cystourethrography, renal scarring, urinary tract infection, reflux nephropathy, continuous antibiotic prophylaxis, endoscopic subureteric injection, dimercaptosuccinic acid scintigraphy, ureteral reimplantation.*

Today, vesicoureteral reflux - defined as the retrograde flow of urine from the urinary bladder into one or both ureters and the renal collecting system - remains among the most clinically consequential urological conditions of childhood. It is identified in approximately 30-40% of children presenting with febrile urinary tract infections and constitutes a leading contributor to pediatric hypertension and chronic kidney disease, with a reported prevalence of 1-2% in the general pediatric population worldwide.

The clinical presentation of vesicoureteral reflux is characteristically heterogeneous, ranging from asymptomatic bacteriuria detected incidentally on routine screening to recurrent febrile urinary tract infections accompanied by

progressive renal parenchymal deterioration. The internationally accepted grading system classifies reflux on a five-point scale: grade I denotes retrograde urine flow into a non-dilated ureter, while grade V represents massive ureteral dilatation with severe tortuosity and intrarenal reflux. This classification system underpins both prognostic stratification and therapeutic decision-making, as spontaneous resolution is markedly more probable in lower-grade disease. The pathophysiological mechanism of renal injury in vesicoureteral reflux involves the direct transmission of elevated intravesical pressure to the renal parenchyma, precipitating focal interstitial fibrosis and glomerulosclerosis, a process collectively designated as reflux nephropathy, which may ultimately culminate in end-stage renal failure if left unrecognized or inadequately managed. Diagnostic confirmation relies principally on voiding cystourethrography, which permits precise anatomical characterization and grading of the refluxing unit. Owing to concerns regarding ionizing radiation exposure inherent to this modality, contrast-enhanced voiding urosonography has emerged as a clinically valid radiation-free alternative increasingly adopted in specialized centers. Assessment of renal cortical integrity is most reliably accomplished through technetium-99m-labeled dimercaptosuccinic acid scintigraphy, a nuclear imaging technique that delineates areas of parenchymal scarring with markedly superior sensitivity relative to conventional renal ultrasonography. Therapeutic management encompasses both conservative and interventional modalities, selected according to reflux grade, patient age, infection history, and renal functional status. Continuous low-dose antibiotic prophylaxis constitutes the cornerstone of conservative management, directed at preventing recurrent urinary tract infections while awaiting spontaneous reflux resolution. Endoscopic subureteric injection of bulking agents - most notably dextranomer/hyaluronic acid copolymer - provides a minimally

invasive surgical alternative with reported success rates of 70-85% in moderate-grade disease. Open surgical ureteral reimplantation, employing the Cohen cross-trigonal or the Lich-Gregoir extravesical technique, remains the definitive gold standard for high-grade or medically refractory reflux, achieving anatomical correction rates exceeding 95%.

In conclusion, effective management of vesicoureteral reflux in children demands individualized, multidisciplinary strategies integrating advanced diagnostic imaging, evidence-based risk stratification, and long-term renal surveillance, with future investigation warranted into genetic predisposition and novel minimally invasive therapeutic modalities.

REFERENCES:

1. Elder J.S., Peters C.A., Arant B.S. et al. Pediatric Vesicoureteral Reflux Guidelines Panel Summary Report on the Management of Primary Vesicoureteral Reflux in Children // Journal of Urology. - 1997. - Vol. 157, No. 5. - P. 1846-1851.
2. Tekgul S., Riedmiller H., Hoebeke P. et al. EAU Guidelines on Vesicoureteral Reflux in Children // European Urology. - 2012. - Vol. 62, No. 3. - P. 534-542.
3. Hoberman A., Greenfield S.P., Mattoo T.K. et al. Antimicrobial Prophylaxis for Children with Vesicoureteral Reflux // New England Journal of Medicine. - 2014. - Vol. 370, No. 25. - P. 2367-2376.
4. Skoog S.J., Peters C.A., Arant B.S. et al. Pediatric Vesicoureteral Reflux Guidelines Panel Summary Report: Clinical Practice Guidelines for Screening Siblings of Children with Vesicoureteral Reflux and Neonates with Prenatal Hydronephrosis // Journal of Urology. - 2010. - Vol. 184, No. 3. - P. 1145-1151.